

		FLUSHING PROTOCOL (0.9% Sodium Chloride)*		Locking Protocol (0.9% Sodium Chloride)*	
		Pre-Infusion	Post-Infusion	Post Lab Draw	Post Normal Saline Flush*
ADULT >66 lbs (30 kg)	PIV	3mL	3mL		3mL
	Midline	3mL	3mL		3mL
	Implanted Port	5mL	10mL	10mL	5mL
	PICC	5mL	10mL	10mL	5mL
	Tunneled Catheter	5mL	10mL	10mL	5mL
	Non-tunneled Catheter	5mL	10mL	10mL	5mL
PEDIATRIC 33 lbs-66 lbs (15 kg-30 kg)	PIV	3mL	3mL		3mL
	Midline	3mL	3mL		3mL
	Implanted Port	5mL	10mL	10mL	3mL
	PICC	5mL	10mL	10mL	3mL
	Tunneled Catheter	5mL	10mL	10mL	3mL
	Non-tunneled Catheter	5mL	10mL	10mL	3mL

For pediatrics less than 33 lbs (15 kg), AICC requires flushing orders to be prescribed by the referring provider.

*0.9% Sodium Chloride will be substituted with Dextrose 5%, or alternative, only when indicated due to medication incompatibility with 0.9% Sodium Chloride. Heparin Sodium can be utilized by provider order.

Advancing quality. Improving lives.